arXiv:1304.7502v1 [math.CV] 28 Apr 2013

GUIDING ISOTOPIES AND HOLOMORPHIC MOTIONS

FREDERICK. P. GARDINER AND YUNPING JIANG

Abstract. We develop a kind of isotopy principle for holomorphic motions. Our main result concerns the extendability of a holomorphic motion of a finite subset E of a Riemann surface Y parameterized by a point t in a pointed hyperbolic surface (X, t_0). It says that if the motion from E to E_t in Y has a guiding quasiconformal isotopy, then an extension to any new point p in $Y-E$ is holomorphically possible. The proof gives a canonical way to replace a motion of the $(n+1)-st$ point by a holomorphic motion while leaving unchanged the given holomorphic motion of the first n points. In particular, our main result gives a new proof of Slodkowski's theorem which concerns the special case when the parameter space is the open unit disk with basepoint 0 and the dynamical space Y is the Riemann sphere.

1. Introduction

Suppose we are given a finite set $\{p_1(t), \ldots, p_n(t)\}$ of holomorphic maps from a hyperbolic pointed Riemann surface (X, t_0) with values in a Riemann surface Y such that the set $E_t = \{p_1(t), \ldots, p_n(t)\}$ consists of n distinct points in Y for every $t \in X$. Here X is called the parameter space and Y the dynamical space. Suppose in addition we are given a continuous function $X \ni t \mapsto p(t) \in Y$ such that for every value of $t \in X$, $E_t \cup \{p(t)\}$ consists of $n+1$ distinct points in Y. If all of these hypotheses are satisfied, then $E_t, t \in X$, is a holomorphic motion in Y with parameter space X which is extended continuously by the motion $E_t \cup \{p(t)\}$ in Y with the same parameter space X. The main goal of this paper is to give an additional condition on $p(t)$ that provides a canonical way to replace the continuous function $p(t) : X \to Y$ with a holomorphic function $\widehat{p}(t) : X \to Y$ such that $\widehat{p}(t) \neq p_i(t)$ for all $t \in X$ and $1 \leq i \leq n$ and such that $p(t_0) = \widehat{p}(t_0)$. This will give us a holomorphic motion $E_t \cup \{\widehat{p}(t)\}$ with parameter space X.

Our main result says that this is possible as long as we are given a guiding isotopy for the continuous motion $E_t \cup \{p(t)\}$. In particular, we give a new

---

2000 *Mathematics Subject Classification.* Primary 30F60; Secondary 32G15, 30C70, 30C75.

The second author is partially supported by the collaboration grant from the Simons Foundation (No.199837) , the CUNY collaborative incentive research grant (No.1861), awards from PSC-CUNY, and a grant from the NSF of China (No.11171121), and a collaboration grant from Academy of Mathematics and Systems Science and the Morningside Center of Mathematics at the Chinese Academy of Sciences.

proof of Slodkowski's theorem which concerns the case when the pointed parameter space (X, t_0) is the open unit disk Δ with base point $t_0 = 0$ and the dynamical Riemann surface Y is the Riemann sphere $\overline{\mathbb{C}}$ minus three points. The proof has a topological part and a geometric part. The topological part says that any holomorphic motion of a finite subset E of the Riemann sphere with parameter space $(\Delta, 0)$ is always guided by a guiding quasiconformal isotopy with the same parameter space. The geometric part assumes there is given a guiding isotopy parameterized by the Riemann surface (X, t_0) for a motion in the Riemann surface Y. Assuming this it provides a canonical holomorphic replacement. It relies on the existence of a canonical cylindrical differential with a double pole at any point, on the heights mapping for quadratic differentials and on the use of harmonic Beltrami differentials to produce holomorphic coordinates for Teichmüller space.

The paper is organized as follows. In section 2, we review definitions of continuous, quasiconformal, and holomorphic motions and introduce the notion of a guiding quasiconformal isotopy. In section 3, we state our main results,Theorems 1 and 2, and Slodkowski's theorem. In section 4, we present the theory of holomorphic quadratic differentials, the heights mapping theorem and a limiting process which yields a cylindrical quadratic differential with a single infinite cylinder. In section 5, we define our extension that follows a given guiding isotopy. In section 6, we prove that this extension is actually a holomorphic motion. The main ideas in sections 5 and 6 concern extremal infinite cyclinders corresponding to quadratic differentials with double poles and on the use of harmonic Beltrami differentials as coordinates for Teichmüller space. In section 7, we give our new proof of Slodkowski's theorem. In section 8 we use Fuchsian groups and their deformations to quasi-Fuchsian groups to generalize the theorem from case where $Y = \overline{\mathbb{C}} - \{0, 1, \infty\}$ to the case that Y is an arbitrary Riemann surface. To make the paper self-contained, we also show how to promote holomorphic motions of finite subsets of Y to holomorphic motions of all of Y. In section 9 we discuss topological obstructions that show that motions with non simply-connected parameter spaces do not necessarily have guiding isotopies. Thus the guiding isotopy assumption is necessary.

Acknowledgement: We would like to thank Linda Keen, Sudeb Mitra, Hiroshige Shiga, and Zhe Wang for helpful discussions.

2. Motions and Guiding Isotopies.

In this section we define motions of a set E in a fixed Riemann surface Y and guiding isotopies of motions. We assume the motion is parameterized by points t in a pointed Riemann surface (X, t_0).

Definition 1. Let $E \subset Y$. A motion h of E in Y parameterized by X with basepoint t_0 is a continuous map $h(t, z) : X \times E \to Y$ satisfying

1) $h(t_0, p) = p$, for all $p \in E$,
2) for any fixed $t \in X$, $E \ni p \mapsto h(t, p) \in Y$ is injective, and
3) for any fixed $p \in E$, $X \ni t \mapsto h(t, p_j) \in Y$ is continuous.

We think of the parameter t as measuring time and $h_t(z) = h(t, z)$ as specifying the motion of the point z in the dynamical space.

Definition 2. A motion h is called a holomorphic motion of E if for each $p \in E$, $X \ni t \to h(t, p) \in Y$ is holomorphic.

Definition 3. Suppose $h(t, z)$ and $\hat{h}(t, z)$ are motions of subsets E and $\hat{E}$ in Y. If $E \subset \hat{E}$ and $\hat{h}(t, z) = h(t, z)$ for all z in E, then $\hat{h}$ is called an extension of h to $\hat{E}$.

Definition 4. A guiding isotopy for a motion h of $E \subset Y$ is an extension $H : X \times Y \to Y$ of h to all of Y. It is called a quasiconformal guiding isotopy if for each $t \in X$, $Y \ni z \to H(t, z) \in Y$ is a quasiconformal.

We have the following proposition about guiding isotopies.

Proposition 1. *Suppose $h(t, z) : X \times E \to Y$ is a holomorphic motion. If h has a quasiconformal guiding isotopy $H(t, z) : X \times Y \to Y$, then any quasiconformal isotopy $G : (t, z) : X \times Y \to Y$ is isotopic to H on $Y - E$.*

Proof. For any $t \in X$, let $H_t = H(t, \cdot)$ and $G_t = H(t, \cdot)$. Then both of them are quasiconformal homeomorphisms of Y. Let $F_t = G_t^{-1} \circ H_t$. Since both of them are extensions of h, we have that $F_t(z) = z$ for all $z \in E$. Since F_t is a quasiconformal homeomorphism, we can define its Beltram coefficient $\mu_t = (F_t)_{\overline{z}}/(F_t)_z$ on Y. It is continuous on $t \in X$. Thus $\phi(t) = F_t$ is a continuous map from X to the space of homeomorphisms of Y. But $\phi(t_0) = F_{t_0} = Id$. $\square$

This proposition shows that the isotopy class of the extension H of h relative to $Y - E$ is unique. However, its isotopy class relative to $Y - E \cup \{p\}$ may not be unique. Our recipe for constructing a holomorphic extension of h to $E \cup \{p\}$ depends only the isotopy class of H relative to $E \cup \{p\}$.

3. Statement of the main results

In the following we let the parameter space be a pointed Riemann surface (X, t_0) and the dynamical space be another hyperbolic Riemann surface Y.

Theorem 1 (Extending motions of finite sets)**.** *Suppose*

$$X \times Y \ni (t,z) \mapsto H(t,z) \in Y$$

is a guiding quasiconformal isotopy for a holomorphic motion $p_j(t) = h(t,p_j)$ *of* n *distinct points*

$$E_t = \{p_1(t), \cdots, p_n(t)\}$$

in Y*. Then there for any point* $p \in Y$ *not equal to any of the points of* $E = \{p_1(t_0), \ldots, p_n(t_0)\}$*, the guiding quasiconformal isotopy* $H(t,z)$ *determines a holomorphic motion* $\hat{h}(t,p)$ *that extends the given holomorphic motion* $h(t,z)$ *of* E *to the set* $E \cup \{p\}$*. The holomorphically moving point* $\hat{h}(t,p)$ *depends only on the guiding isotopy class of* H *relative to* $E \cup \{p\}$*.*

The complete proof of this theorem is given at the end of section 8.

The next theorem can be obtained from this one by letting Y equal $\mathbb{C}_{0,1}$, that is, the Riemann sphere with $\{0,1,\infty\}$ removed. However, our approach will be the opposite. First we prove the next theorem which treats the case when Y is the complement in the Riemann sphere of a set of three or more points. Then we realize the surfaces Y, Y_ϵ and $Y_{t,\epsilon}$ that come up in the proof by Fuchsian and Kleinian groups that act on the sphere and, by this means, we obtain a proof of Theorem 1 derived from the the proof of Theorem 2.

Suppose $p_j(t) = h(t,p_j)$ is a continuous motion of $E = \{p_1, \ldots, p_n\} \subset \overline{\mathbb{C}}$. Since the points of E move continuously and distinctly, there is a continuous path of Möbius transformations A_t such that $A_t(p_1(t) = 0, A_t(p_2(t)) = 1$ and $A_t(p_3(t)) = \infty$. Then the ordered n-tuple

$$A_t(E_t)) = (0, 1, \infty, A_t(p_4(t)), \ldots, A_t(p_n(t)))$$

is a continuous motion of $A_0(E)$. Obviously, if E_t depends holomorphically on t then so does $A_t(E_t)$. Thus by showing how to extend a holomorphic motion of a subset of $\overline{\mathbb{C}}$ normalized at $0, 1, \infty$, by choosing an appropriate holomorphic path of Möbius transformations A_t we also provide a method for extending holomorphic motions that are not normalized.

Theorem 2 (Motions in the Riemann sphere)**.** *Suppose*

$$E = \{p_1, \cdots, p_n\}$$

is a finite subset of $\overline{\mathbb{C}}$ *containing* $n \geq 3$ *distinct points and* X *is a hyperbolic Riemann surface with base point* t_0*. Assume* $h : X \times E \to \overline{\mathbb{C}}$ *is a holomorphic motion of* E *in* $\overline{\mathbb{C}}$ *and* p *is a point in* $\overline{\mathbb{C}} - E$*. If there is a guiding quasiconformal motion* $\widetilde{h} : X \times \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ *that extends* h*, then there is a holomorphic motion* $\widehat{h} : X \times (E \cup \{p\}) \to \overline{\mathbb{C}}$ *that has the same initial value at* $t = t_0$ *as* $\widetilde{h}$ *and that extends* h*. The holomorphically moving point* $\widehat{h}(t,p)$ *depends only on the guiding isotopy class of* $\widetilde{h}$ *relative to* $E \cup \{p\}$*.*

If the parameter space X is compact or compact except for a finite number of punctures, this theorem has no interest because the only holomorphic functions are constant.

The next sections up to and including section 7 are devoted to proving Theorem 2. In section 8 we show how to use Theorem 2 to obtain Theorem 1. The proofs rely on properties of holomorphic quadratic differentials with double poles, see [24], and on holomorphic coordinates for Teichmüller space provided by the Bers' embedding, see [4].

Letting (X, t_0) be the unit disc Δ with base point 0, Theorem 2 becomes the following theorem of Slodkowski.

Theorem 3 (Slodkowski [23]). *Suppose* $X = \Delta = \{t \mid |t| < 1\}$ *is the open unit disk, with the base point* $t_0 = 0$. *Then any holomorphic motion* $h(t, z) : \Delta \times E \to \overline{\mathbb{C}}$ *can be extended to a holomorphic motion* $\widetilde{h}(t, z) : \Delta \times \overline{\mathbb{C}} \to \overline{\mathbb{C}}$.

4. CYLINDERS WITH MAXIMAL MODULUS

Theorem 4 (Heights Mapping Theorem). *Assume* E *is a finite subset of* $\overline{\mathbb{C}}$ *containing two or more points and let* $\Delta(p)$ *be a conformal disc centered at* p *in* $\overline{\mathbb{C}} - E$. *Let* γ *be a simple closed curve in* $Y = \overline{\mathbb{C}} - (E \cup \Delta(p))$ *and homotopic to the boundary of* $\Delta(p)$. *Then there exists a unique holomorphic quadratic differential* q *of finite norm defined on* Y *with the following properties:*

1. *all of the regular horizontal trajectories of* q *are closed and homotopic to* γ,
2. *the regular horizontal trajectories form a cylinder* A *that fills* $\overline{\mathbb{C}} - (E \cup \Delta(p))$ *except for a critical graph* C,
3. *each regular horizontal trajectory* α *in this cylinder has length equal to* 2π *and the totality of these trajectories fill* A,
4. C *is closed, connected set of measure zero and is the union of critical horizontal trajectories of* q *that join its zeros and poles,*
5. C *is a connected, simply connected finite graph, the poles of* q *form the endpoints of* C *and any zero of order* k *is a vertex of* C *from which emanates* $k + 2$ *edges of* C, *and*
6. $||q|| = \int\int_{\overline{\mathbb{C}} - (E \cup \Delta(p))} |q(z)| dxdy = 2\pi b$, *where* b *is the height of* A *measured in the metric* $|q|^{1/2}$.

q *is the unique holomorphic quadratic differential on* Y *with the properties that it has a characteristic cylinder with maximal modulus among all cylinders homotopic to the boundary of* $\Delta(p)$ *and its circumference measured with respect to the metric* $|q(z)|^{1/2}|dz|$ *is equal to* 2π.

Proof. The existence and uniqueness of q with these properties is well-known, see [9, 12, 16, 24]. For any simple closed curve α not homotopically trivial and not homotopic to a puncture on any hyperbolic Riemann surface, such a holomorphic quadratic differential is obtained by maximizing the modulus of a cylinder among all cylinders on the surface homotopic to α. A similar conclusion is true even if we begin with a system $\{\alpha_j\}$ of non-homotopic simple closed curves, and either the heights of the annuli can be arbitrarily specified (the heights theorem of Renelt [21]) or the projective class of the moduli of the annuli can be arbitrarily specified (the Strebel moduli theorem [24]).

For completeness of exposition here we give a sketch of the proof in the case we need, namely, the case where there is just one annulus with core curve homotopic to a boundary component and the size of this boundary component shrinking to zero.

Lemma 1. *Suppose A is an annular Riemann surface conformal to the ring domain $\{z : r < |z| < R\}$ and $\mu(z)$ is an L^∞ Beltrami differential on A with $||\mu||_\infty < 1$. In terms of the conformal parameter w let $q(w)(dw)^2 = \left(\frac{dz}{z}\right)^2$ and let A_μ be the same annulus with the conformal structure induced by μ. This means a local homeomorphism $w = h(z)$ from a neighborhood of a point p in A is declared to be conformal if*

$$h_{\overline{z}}(z) = \mu(z)h_z(z).$$

Let $\Lambda(A)$ be the extremal length of the family of arcs in A that join its two boundary components and $\Lambda(A_\mu)$ be the extremal length with the same family with respect to the conformal structure induced by μ. Then

$$\log \Lambda(A_{t\mu}) = \log \Lambda(A) + 2 \text{ Re } \frac{t}{||q||} \iint_A \mu q dxdy + O(|t|^2). \tag{1}$$

Proof. This formula follows from the Reich-Strebel inequalities and is proved in [10] and [13]. □

Lemma 2. *Suppose A is an annulus embedded in a Riemann surface Y and the modulus of A is a large as possible among all annuli homotopic to A in Y. Let g be a conformal map from A onto the region $\{z : r < |z| < R\}$. Then $-\left(\frac{dg}{g}\right)^2$ is the restriction of a holomorphic quadratic differential q on Y and*

$$\iint_Y |q|dxdy = \iint_A \left|\frac{1}{z^2}\right| dxdy = 2\pi \log(R/r).$$

All of the regular horizontal trajectories of q are closed and are the images under g of the circles $\rho e^{i\theta}, r < \rho < R$.

Proof. This lemma is proved in [9] and [12]. For the benefit of the reader we repeat the main ideas of the proof here.

The complex Banach space $Q(Y)$ of integrable holomorphic quadratic differentials on Y is a closed subspace of $L^1(Y)$, the space of integrable quadratic differentials and the dual Banach space $L^1(Y)^*$ is isometric to L^∞, the space of essentially bounded Beltrami differentials under the pairing

$$(q, \mu) = \iint_Y \mu q dxdy.$$

In particular, if μ represents a linear functional $\ell \in L^1(Y)^*$ then

$$||\ell||_* = \sup_{q \in L^1(Y)} \frac{\left(\iint \mu q dxdy\right)}{||q||} = ||\mu||_\infty .$$

Our strategy to prove this lemma uses the identification of $Q(Y)^{\perp\perp}$ with $Q(Y)$ provided by this pairing. Showing that $\left(\frac{dg}{g}\right)^2$ is in $Q(Y)^{\perp\perp}$ also shows that it is in $Q(Y)$. If $\left(\frac{dg}{g}\right)^2 \notin Q(Y)^{\perp\perp}$ then there is a Beltrami differential μ supported in A such that

$$\iint_A \mu q dxdy = 0$$

for all $q \in Q(Y)$ and $\iint_A \mu \left(\frac{dg}{g}\right)^2 dxdy \neq 0$. By the Hamilton-Krushkal variational lemma there would be a curve of deformations μ_t tangent to μ at $t = 0$ for which by Lemma 1 $\Lambda(A_{\mu_t})$ is smaller than $\Lambda(A)$ and for which $Y(\mu_t)$ represents the same point of Teichmüller space as Y. This conclusion contradicts the assumption that $\Lambda(A)$ has maximum modulus among homotopic annuli A embedded in Y. ☐

The following result is also well known [12, 14].

Lemma 3. *Let f be a quasiconformal homeomorphism mapping a hyperbolic Riemann surface Y to a Riemann surface $f(Y)$ and let K be the maximal dilatation of f. Let q be a holomorphic quadratic differential on Y of finite norm with given heights and q_f a holomorphic quadratic differential on $f(Y)$ such the height along the isotopy class of any curve γ in Y measured with respect to $|\mathrm{Im}\ (q(z)^{1/2}dz)|$ is equal to the height of the isotopy class $f(\gamma)$ in $f(Y)$ measured with respect to $|\mathrm{Im}\ (q_f(w)^{1/2}dw)|$. Then*

$$K^{-1}||q|| \leq ||q_f|| \leq K||q||. \tag{2}$$

Proof. By the Dirichlet principle [11, 12] for measured foliations

$$||q|| = \iint_Y |q| dxdy$$

is equal to the infimum of $2D(v) = 2\int\int_Y (v_x^2 + v_y^2)\ dxdy$, where the infimum is taken over all measured foliations $|dv|$ that realize the heights of q, and any measured foliation that realizes this infimum is the absolute value of the imaginary part of the square root of q. Note that $|d(v \circ f)|$ has the same corresponding heights on Y that $|dv|$ has on $f(Y)$, and

$$(v \circ f)_z = (v_w \circ f) f_z + (v_{\overline{w}} \circ f)\overline{f}_z .$$

Since v is real-valued and defined up to plus or minus sign and up to the addition of a constant, $|v_w|$ and $|v_{\overline{w}}|$ are invariant and $|v_w|^2 = |v_{\overline{w}}|^2 = (1/4)(v_x^2 + v_y^2)$. Thus we can use exactly the same calculation that is given at the end of Chapter 1 in [1]. We have

$$|(v \circ f)_z| \leq (|v_w| \circ f)(|f_z| + |f_{\overline{z}}|), \text{ and}$$

$$D_Y(|dv \circ f|) = 2 \iint_Y |(v \circ f)_z|^2 |dz \wedge \overline{dz}| \leq$$

$$2 \iint_Y (|v_w| \circ f)^2 (|f_z| + |f_{\overline{z}}|)^2 |dz \wedge d\overline{z}| =$$

$$2 \iint_{f(Y)} |v_w|^2 \ \frac{(|f_z| + |f_{\overline{z}}|)^2}{|f_z|^2 - |f_{\overline{z}}|^2} \ |dw \wedge d\overline{w}| =$$

$$2 \iint_{f(Y)} |v_w|^2 \left(\frac{|f_z| + |f_{\overline{z}}|}{|f_z| - |f_{\overline{z}}|} \right) |dw \wedge d\overline{w}| \leq K(f) D_{f(Y)}(|dv|).$$

This implies the left hand side of (2). The right hand side follows from the same argument applied to f^{-1}. $\square$

Lemmas 1, 2, and 3 complete the proof of Theorem 4. $\square$

5. Definition of the extension

In this section we assume we are given a holomorphic motion $E_t = \{p_1(t), \ldots, p_n(t)\}$ of a finite set $E = \{p_1, \ldots, p_n\}$ in $\overline{\mathbb{C}}$ parameterized by $t \in X$. Without loss of generality, we always assume that $p_1(t) = 0, p_2(t) = 1$ and $p_3(t) = \infty$ for every $t \in X$. Suppose in addition to the holomorphic functions $p_j(t)$, $1 \leq j \leq n$, defined for $t \in X$, we are given another continuous function $p(t)$ in $\overline{\mathbb{C}}$ so that $E_t \cup \{p(t)\} = \{p_1(t), \ldots, p_n(t), p(t)\}$ is a set of $n+1$ distinct points for each $t \in X$. If we let

$$h(t,z) = \begin{cases} p_j(t) & \text{for } z = p_j(t_0), 1 \leq j \leq n \text{ and} \\ p(t) & \text{for } z = p(t_0), \end{cases}$$

then h is a motion, which is holomorphic on E and continuous on $p = p(t_0)$. Now select a disc $\Delta(p, \epsilon)$ of radius ϵ centered at p where ϵ is less than the shortest distance $|p(t_0) - p_j(t_0)|, 1 \leq j \leq n$. Let $G(t,z) = \widetilde{h}(t,z)$ be the guiding isotopy in the assumption of Theorem 2 such that $p(t) = G(t,p)$. Because each map $z \mapsto G(t,z)$ is quasiconformal, $\Delta(t, \epsilon) = G(t, \Delta(p, \epsilon))$ is a quasidisc disjoint from E_t for each $t \in X$. Our goal is to use G to define a replacement $\hat{h}$ of $G|(E \cup \{p\})$ by means of a limiting process in such a way that $\hat{h}$ is a holomorphic motion of $E \cup \{p\}$ that extends the holomorphic motion of E given by h. The extension $\hat{h}$ of h will depend on the equivalence class relative to $E \cup \{p\}$ of the guiding quasiconformal isotopy G. Also, we want $\hat{h}$ to have the same initial point as G and in some sense we want it to follow a path topologically equivalent to the path $t \mapsto p(t)$. We make the definition of $\hat{h}(p)$ in several steps.

Step I. For $w \in \mathbb{C}$ let $\Delta(p,\epsilon) = \{w : |w-p| < \epsilon\}$ and let j be the reflection across the boundary of $\Delta(p,\epsilon)$ defined by

$$j(w) = p + \epsilon^2/(\overline{w-p}). \tag{3}$$

Let

$$Y_\epsilon = \overline{\mathbb{C}} - (E \cup j(E)) \, .$$

By Theorem 4 we construct the cylinder A_ϵ in $Y_\epsilon = \overline{\mathbb{C}} - (E \cup j(E))$ with maximal modulus and with core curve homotopic to the boundary of $\Delta(p,\epsilon)$. It determines a holomorphic quadratic differential q_ϵ with characteristic cylinder A_ϵ and a conformal map $w \mapsto z$ from that cylinder onto a domain in the plane of the form

$$\{z : 1 < |z| < r\},$$

where $(1/2\pi)\log r$ is the modulus of the cylinder. In this annulus

$$q_\epsilon(w)(dw)^2 = -\left(\frac{dz}{z}\right)^2$$

and because of the uniqueness of q_ϵ under the symmetry j, the boundary of $\Delta(p,\epsilon)$ is a regular horizontal trajectory of q_ϵ. Note that the domain $Y_\epsilon = Y - (E \cup j(E))$ is normalized by the condition that the first three points of E are $0, 1$ and ∞. The horizontal foliation with its vertical measure is given by

$$|d\theta| = |\mathrm{Im}\ \left(((-q_\epsilon)^{1/2})(w)dw\right)| = |\mathrm{Re}\ (q_\epsilon(w)(dw)^2)^{1/2}|.$$

Step II. Let $\tilde{\mu}_{t,\epsilon}$ be the Beltrami coefficient of $z \mapsto G_t(z) = G(t,z)$, let

$$\mu_{t,\epsilon} = \begin{cases} 0 & \text{in } \Delta(p,\epsilon) \\ \tilde{\mu}_{t,\epsilon} & \text{in } \overline{\mathbb{C}} - \Delta(p,\epsilon), \end{cases} \tag{4}$$

and let $g^{t,\epsilon}$ be the unique quasiconformal self-mapping of $\overline{\mathbb{C}}$ that fixes $0, 1$ and ∞ and that has Beltrami coefficient $\mu_{t,\epsilon}$.

Step III. For each t, the isotopy type of the quasiconformal map $z \mapsto g^{t,\epsilon}(z)$ determines a heights mapping from Y_ϵ to $Y_{t,\epsilon}$. Here we define the quasidisc $\Delta(t,\epsilon)$ by the equation

$$\Delta(t,\epsilon) = g^{t,\epsilon}(\Delta(p,\epsilon)), \tag{5}$$

and we put $Y_{t,\epsilon}$ equal to $g^{t,\epsilon}(Y_\epsilon)$. By the heights mapping theorem, Theorem 4, there is a unique holomorphic quadratic differential $q_{t,\epsilon}$ on $Y_{t,\epsilon}$ such that the heights of

$$|d\theta_t| = |\mathrm{Re}\ (q_{t,\epsilon}(w)(dw)^2)^{1/2}|$$

on $Y_{t,\epsilon}$ are equal to the corresponding heights of $|d\theta| = |\mathrm{Re}\ (q_\epsilon(w)(dw)^2)^{1/2}|$ on Y_ϵ. In particular the height of $|d\theta_t|$ along the boundary of $\Delta(t,\epsilon)$ is equal to 2π.

Note that the critical graph of q_ϵ consists of two trees, the first of which has endpoints at the points of E and the second of which has endpoints at the points of $j(E)$. Similarly, the critical graph of $q_{t,\epsilon}$ also consists of two trees and, correspondingly, the first has endpoints at the points of $g^{t,\epsilon}(E)$ and the second has endpoints at the points of $g^{t,\epsilon}(j(E))$.

Step IV. By the heights mapping theorem and the minimum Dirichlet principle for measured foliations, [11, 15],

$$(1/K_t)\iint_{\overline{\mathbb{C}}}|q_\epsilon| \le \iint_{\overline{\mathbb{C}}}|q_{t,\epsilon}| \le K_t\iint_{\overline{\mathbb{C}}}|q_\epsilon|. \tag{6}$$

The family of simple closed curves that are homotopic to the boundary of the quasidisc $\Delta(t,\epsilon)$ in $\overline{\mathbb{C}} - g^{t,\epsilon}(E \cup j(E))$ has extremal length less than

$$2\pi/\iint_{\overline{\mathbb{C}}}|q_{t,\epsilon}|.$$

Since $\int\int_{\overline{\mathbb{C}}}|q_\epsilon| \to \infty$ as $\epsilon \to 0$, by holding t fixed and letting ϵ approach zero, we see that the domain $g^{t,\epsilon}(A_\epsilon)$ approaches a domain in the Riemann sphere conformal to a punctured disc. All of the points of $g^t(j(E))$ are squeezed to this puncture. Moreover, since $g^{t,\epsilon}$ is Hölder continuous in the sense that there is a constant C such that

$$|g^{t,\epsilon}(w_1) - g^{t,\epsilon}(w_2)| \le C|w_1 - w_2|^{1/K_t},$$

every point of the closure of $\Delta(t,\epsilon) = g^{t,\epsilon}(\Delta(p,\epsilon))$ converges to this puncture as $\epsilon \to 0$.

Definition 5. We put

$$\hat{h}(t,z) = \begin{cases} \hat{p}(t) & \text{for } z = p(t_0) \text{ and} \\ \lim_{\epsilon\to 0} g^{t,\epsilon}(z) & \text{for } z = p_j(t_0), 1 \le j \le n. \end{cases} \tag{7}$$

Theorem 5. *The function $\hat{h}(t,z)$ is a motion of $E \cup \{p\}$ that extends the holomorphic motion h of E. Moreover, for no value of $t \in X$ can $\hat{h}(t,p)$ belong to E_t because it is separated from E_t by annuli of arbitrarily large moduli. This means that for each t, the extension $\hat{h}$ of h is injective.*

Proof. The map $g^{t,\epsilon}$ defined in Step II depends on the reflection defined in Step I, which depends on ϵ. As $\epsilon \to 0$, the Beltrami coefficients $\mu_{t,\epsilon}$ converge in the bounded pointwise sense to the Beltrami coefficient of $z \mapsto G(t,z)$ and so since these mappings are normalized to fix $0, 1$ and ∞, $g^{t,\epsilon}$ converges uniformly in the spherical metric to $z \mapsto G(t,z)$. Therefore, $\hat{h}(p_j(t)) = h(p_j(t))$ for $p_j(t) \in E_t$. Since $\hat{h}(p(t))$ is separated from all of the points of E_t by an annulus of positive modulus, it cannot coincide with any of those points.

The limiting differential $q_t = \lim_{\epsilon\to 0} q_{t,\epsilon}$ is a holomorphic on the Riemann sphere except for possibly simple poles at E_t and a double pole with quadratic residue equal to -1 at $\hat{h}(t,p)$. Since its construction depends on the guiding isotopy G, in the remaining part of this proof, we denote $q_{t,\epsilon}$ by $q_{t,\epsilon,G}$. □

In the next section we will show that $\hat{h}$ is also a holomorphic extension of h, that is, there is a neighborhood N of every value of $s_0 \in X$ such that for $t \in N$, $\hat{h}(t, z)$ depends holomorphically on t for $z \in E \cup \{p\}$. We will show that $\hat{p}(t)$ depends holomorphically on t by showing that the points for fixed $\epsilon > 0$ the points $g^{t,\epsilon}(z)$ where z is any point in the closure of $\Delta(t, \epsilon)$ depend holomorphically on the holomorphic functions $p_j(t), 1 \leq j \leq n$, and for any such z, $g^{t,\epsilon}(z)$ converges uniformly to $\hat{h}(t, p)$ as $\epsilon \to 0$.

6. HARMONIC COORDINATES FOR TEICHMÜLLER SPACE

In this section we use harmonic Beltrami coefficients as coordinates for the unreduced Teichmüller space of the bordered Riemann surface $R = Y_\epsilon - \Delta(p, \epsilon)$. This Teichmüller space consists of quasiconformal maps f from R to $f(R)$ factored by an equivalence relation. Two such maps f_0 and f_1 are equivalent if, after postcomposing one of them by a conformal map, they can be connected by a isotopy g_t which pins down the points of E and all of the points on the boundary of $\Delta(p, \epsilon)$. In particular, the Teichmüller space is infinite dimensional and its coordinates at the point $[f]$ keep track of the boundary curve of R, namely, the boundary of the quasidisc $f(\Delta(p, \epsilon))$.

From the Ahlfors-Weill extension procedure (see [2]) about the existence of a local holomorphic section, a local coordinate at a point $\tau = [f]$ in $T(R)$ is given by harmonic Beltrami coeffients μ on $f(R)$, which have the following special properties:

1) μ is identically equal to zero on $f(\Delta(p, \epsilon))$,

2) $\mu = \rho^{-2}(z)\overline{\psi(z)}$ where ψ is a holomorphic quadratic differential on $f(R)$ where ρ is the Poincaré metric for $f(R)$, and

3) where $||\mu||_\infty < 1$.

In particular, the holomorphic function $\psi(z)$ has at most simple poles at the points of $f(E)$ and 3) means that $|\psi(z)| \leq C_\psi \delta^{-2}(z)$ where $\delta(z)$ is the minimum Euclidean distance from z to the boundary of $f(\Delta(p, \epsilon))$.

Theorem 6. *Assuming each of the functions $p_j(t), 1 \leq j \leq n$ is holomorphic, the motion $\hat{h}(p, t)$ defined in (7) holomorphically extends the motion $h(p, t)$. In particular,*

a) *$\hat{h}(t, z)$ coincides with $h(t, z)$ for each $z = p_j(t), 1 \leq j \leq n$,*

b) *$\hat{p}(t) = \hat{h}(t, p)$ is distinct from every point of E_t in $\overline{\mathbb{C}}$ for each t in X, and*

c) *$\hat{p}(t)$ is a holomorphic function of $t \in X$.*

Proof. a) was already proved in the previous section.

b) was also noted in the previous section and follows because $\hat{h}(t, p)$ is separated from the points of E_t in $\overline{\mathbb{C}}$ by an annulus of positive modulus.

c) follows from the idea behind the Bers' embedding. First assume the hyperbolic disk $\{x \in X \mid \rho_X(t, t_0) < \delta_0\}$ defines an embedded disc in X where

δ_0 is small enough so that the corresponding points Teichmüller space lie inside a metric ball that has harmonic Beltrami coordinates. In this case since the points $p_j(t), 1 \leq j \leq n$ move holomorphically the harmonic Beltrami coefficients that represent $\mu_{t,\epsilon}$ in the Teichmüller space of $Y_\epsilon - \Delta(p,\epsilon)$ also vary holomorphically. Since these Beltrami coefficients are identically equal to zero inside the disc $\Delta(p,\epsilon)$, all of the points inside this disc vary holomorphically. Therefore, the limiting point $\hat{p}(t)$ as $\epsilon \to 0$ also varies holomorphically since it is the uniform limit as $\epsilon \to 0$ of $g^{t,\epsilon}(p)$, where $g^{t,\epsilon}$ is defined in Step II of the previous section.

Now follow the motion $\hat{h}(t,p)$ to a value $t = s_0$ and apply the same argument to motion starting at E_{s_0} and pick a small enough neighborhood of s_0 where there are harmonic Beltrami coordinates. Then s is a holomorphic function of t and if t is near enough s_0 so that the holomorphically moving equivalence class in $T(Y_\epsilon - \Delta(p,\epsilon))$ is represented by a harmonic Beltrami coefficient μ_s with $||\mu_s||_\infty < 1$. For each $\epsilon > 0$, all of the points of the quasidisc $g^{t,\epsilon}(\Delta(p,\epsilon))$ move holomorphically with respect to t. Thus, we find that the point $\hat{p}(t)$ is a uniform limit of the holomorphic functions $g^{t,\epsilon}(p)$ as $\epsilon \to 0$. Thus $\hat{p}(t)$ is also holomorphic function at every point t with X.

$\square$

7. A new proof of Slodkowski's Theorem

In this section, we give a new proof of Theorem 3. It is based on two results, the first a known topological result and the second geometric result, which is our main Theorem 2. Therefore, to complete our new proof of Theorem 3, we only need to prove the topological result here by showing that the topological condition in Theorem 2 holds when the parameter space X is the open unit disk Δ. More precisely, we prove the following lemma.

Lemma 4. *Suppose Δ is the open unit disk with the basepoint 0. Suppose $E = \{p_1 = 0, p_2 = 1, p_3 = \infty, p_4, \ldots, p_n\}$, $\#(E) = n \geq 3$, is a finite subset of the Riemann sphere $\overline{\mathbb{C}}$ and $h(t,z) : \Delta \times E \to \overline{\mathbb{C}}$ is a normalized holomorphic motion of E. Then h can be extended to a normalized quasiconformal motion $H(t,z) : \Delta \times \overline{\mathbb{C}} \to \overline{\mathbb{C}}$.*

Proof. Let $\Omega = \overline{\mathbb{C}} \setminus E$ be the Riemann sphere punctured at the points of E. The Teichmüller space $T(E)$ is the classical Teichmüller space of the plane domain Ω. From a theorem of Earle and McMullen [8], two quasicconformal deformations w^μ and w^ν of Ω are Teichmüller equivalent if they are connected by a quasiconformal isotopy rel E, where w^μ and w^ν mean the normalized quasconformal homeomorphism with Beltrami coefficients μ and ν.

Let

$$Y_{n-3} = \{\mathbf{z} = (z_1, \cdots, z_{n-3}) \in \mathbb{C}^{n-3}\}$$

where $z_i \neq z_j$ for all $1 \leq i \neq j \leq n-3$ and $z_i \neq 0, 1, \infty$ for all $1 \leq i \leq n-3$. From the normalized holomorphic motion $h(t,z) : \Delta \times E \to \overline{\mathbb{C}}$, we can define a holomorphic map

$$f(t) = (h(t, p_4), \cdots, h(t, p_n)) : \Delta \to Y_{n-3}.$$

We need the following proposition which proof can be found in [5] (see also [20]).

Proposition 2. *The map*

$$\pi_E([\mu]_E) = (w^{\mu}(p_4), \cdots, w^{\mu}(p_n)) : T(E) \to Y_{n-3}$$

is a holomorphic universal covering.

Since Δ is simply connected, we can lift f to get a holomorphic map

$$\widetilde{f}(t) : \Delta \to T(E)$$

such that

$$\pi_E \circ \widetilde{f} = f.$$

From the Douady-Earle barycentric extension procedure (see [6]), there is a continuous section S of P_E (see [17]), that is, a continuous map S from $T(E)$ to $M(\mathbb{C})$ such that $P_E \circ S$ is the identity on $T(E)$. Define

$$\widehat{f}(t) = S \circ \widetilde{f}(t) : \Delta \to M(\mathbb{C})$$

Then we have that

$$P_E \circ \widehat{f} = \widetilde{f} \quad \text{and} \quad \pi_E \circ P_E \circ \widehat{f} = f.$$

Refer to the following diagram:

$$\begin{tikzcd}
 & M(\mathbb{C}) \arrow[d, "P_E"'] \\
 & T(E) \arrow[u, bend right, "S"'] \arrow[d, "\pi_E"] \\
\Delta \arrow[uur, "\widehat{f}"] \arrow[ur, "\widetilde{f}"] \arrow[r, "f"] & Y_{n-3}
\end{tikzcd}$$

For any $\widehat{f}(t) \in M(\mathbb{C})$, let $w^{\widehat{f}(t)}$ be the normalized quasiconformal homeomorphism solving the Beltrami equation

$$w_{\overline{z}} = \mu(z) w_z \tag{8}$$

with the Beltrami coefficient $\mu = \widehat{f}(t)$. Define

$$H(t, z) = w^{\widehat{f}(t)}(z) : \Delta \times \overline{\mathbb{C}} \to \overline{\mathbb{C}}.$$

Then it is a normalized quasiconformal motion of $\overline{\mathbb{C}}$ with parameter space Δ. Since $\pi_E \circ P_E(\widetilde{f}) = f$, we get $H(t, 0) = 0$, $H(t, 1) = 1$, $H(t, \infty) = \infty$, and

$$(H(t, p_4), \cdots, H(t, p_n)) = f(t).$$

Thus H is an extension of h. This completes the proof. $\square$

Lemma 4 says that for any normalized holomorphic motion $h : \Delta \times E \to \overline{\mathbb{C}}$ of any finite subset E of the Riemann sphere $\overline{\mathbb{C}}$ with parameter space Δ, our guiding isotopy assumption in Theorem 2 holds. Thus for any new point $p \notin E$, we can have a holomorphic motion $\widehat{h} : \Delta \times (E \cup \{p\}) \to \overline{\mathbb{C}}$ extending h. To complete the proof, we need the λ-Lemma of Màñé, Sad and Sullivan, (see [19]).

Lemma 5 (λ-Lemma). *Suppose $h(t, z) : \Delta \times E \to \overline{\mathbb{C}}$ is a holomorphic motion, where E is a subset of $\overline{\mathbb{C}}$. Then it can be extended to a holomorphic motion $\overline{h}(t, z) : \Delta \times \overline{E} \to \overline{\mathbb{C}}$, where $\overline{E}$ means the closure of E in $\overline{\mathbb{C}}$.*

Now suppose $h(t, z) : \Delta \times E \to \overline{\mathbb{C}}$ is the normalized holomorphic motion in Theorem 3. Let $E_\infty = \{0, 1, \infty, p_1, \cdots, p_n, \cdots\}$ be a countable dense subset of E. Let $F = \{q_1, \cdots, q_n, \cdots\}$ be a countable dense subset of $\overline{\mathbb{C}} \setminus E$. Let $E_n = \{0, 1, \infty, p_1, \cdots, p_n\}$ and $F_n = \{q_1, \cdots, q_n\}$. Then $h_n = h|\Delta \times E_n$ is a holomorphic motion for every $n > 3$. Our main result (Theorem 2) with the consideration of Lemma 4 implies that we can extend h_n to a holomorphic motion $H_n(t, z) : \Delta \times (E_n \cup F_n) \to \overline{\mathbb{C}}$. Inductively, we have a holomorphic motion $H_\infty(t, z) : \Delta \times (E_\infty \cup F) \to \overline{\mathbb{C}}$ which extends every h_n. The λ-Lemma implies we can extend this last holomorphic motion into a holomorphic motion $H(t, z)$ of the closure of $E_\infty \cup F$ which is the whole Riemann sphere $\overline{\mathbb{C}}$ with parameter space Δ. This holomorphic motion H is an extension of h. This completes our new proof of Theorem 3.

8. Arbitrary dynamical spaces

In this section we show how the proof given in the previous sections for the case when the dynamical space is the Riemann sphere can be promoted to prove the analogous theorem when the dynamical space is any Riemann surface. This is the content of Theorem 1.

We assume that the finite subset E of Y is moving holomorphically and parameterized by a guiding quasiconformal isotopy with parameter $t \in X$. Our goal is to use the isotopy to induce a holomorphically moving point $p(t)$ that starts at any point $p = p(t_0)$ in $Y - E$ and has the property that $p(t)$ never lies in the set E_t.

First, take a point $p \in Y$ and form a Fuchsian universal covering group Γ of $Y - \{p\}$ acting on the upper half plane $\mathbb{H}$ that contains a primitive parabolic element $\gamma_0(z) = z + 1$ with the property that the arc $\tilde{\alpha}(t) = t + iM, 0 \leq t \leq 1$ has image α under the covering that winds once around the point p. By the Leutbecher-Shimizu [18,22] lemma the restriction of the covering mapping to the semi-infinite strip

$$\{w : 0 \leq u = \text{Re } w < 1, v = \text{Im } w > 1\}$$

is a homeomorphism onto an embedded punctured disc contained in $Y - \{p\}$ with the puncture located at p. Thus, $z = \exp(2\pi i w)$ is a local parameter on Y with $z = 0$ corresponding to the point p and the part of this strip lying above $v = M > 1$ corresponding to an embedded disc

$$\{z : |z| \leq \epsilon = \exp(-2\pi M)\}. \tag{9}$$

Now we can replace Step I of section 6 by the following device. We construct a Kleinian group Γ_ϵ that contains two isomorphic copies Γ both of which act discontinuously on a certain domain. The domain has an anticonformal involution j and the action of one copy of Γ covers Y_ϵ and the action of $j \circ \Gamma \circ j$ on the other copy covers $j(Y_\epsilon)$. It turns out that Γ_ϵ is called an HNN-extension of Γ, named after Higman, Neumann and Neumann.

To construct Γ_ϵ, we take $M > 1$ and let j be reflection in the plane around the horizontal line $y = M$. Then we form Kleinian group Γ^d generated by Γ and $j \circ \Gamma \circ j$. Note that both of these groups contain the common element $z \mapsto z + 1$. If we take a fundamental domain for Γ that is bounded by the two vertical lines $x = 0$ and $x = 1$ and axes of hyperbolic transformations all lying below the line $y = 1$, then Γ^d has a fundamental domain lying between the two horizontal lines $y = 0$ and $y = 2M$. This rectangle contains the subrectangle $0 \leq x \leq 1$ and $1 \leq y \leq 2M - 1$, which is entirely contained within the fundamental domain for Γ^d. Moreover, the fundamental domain is invariant under its reflection j across the line $y = M$. Γ^d is the amalgamated product of Γ and $j \circ \Gamma \circ j$, amalgamated along the common cyclic subgroup generated by $z \mapsto z + 1$. Since Γ^d depends on M and M depends on ϵ by the simple formula (9), we denote Γ^d by Γ_ϵ. The groups Γ and Γ_ϵ both act on the Riemann sphere and, if we hold the limit points $0, 1$, and ∞ fixed and let $\epsilon \to 0$, the fundamental domains for the groups Γ_ϵ converge in the Hausdorff sense to the fundamental domain for Γ.



Figure 1

Steps II, III and IV can be repeated almost verbatim but now with Beltrami coefficients $\tilde{\mu}_{t,\epsilon}$ and $\mu_{t,\epsilon}$ and corresponding mappings g^t and $g^{t,\epsilon}$ all equivariant for the Kleinian groups Γ_ϵ. As one lets $M \to \infty$ or $\epsilon \to 0$ and holds three limit points $0, 1$ and ∞ fixed, the groups Γ_ϵ converge to Γ, and so do their corresponding conjugated groups $g^{t,\epsilon} \circ \Gamma_\epsilon \circ (g^{t,\epsilon})^{-1}$ converge to $g^t \circ \Gamma \circ (g^t)^{-1}$. The extension of the holomorphic motion to p is realized by the corresponding fixed point of a parabolic transformaton in $g^t \circ \Gamma \circ (g^t)^{-1}$. Since the entries of this parabolic transformation vary holomorphically with t, so does the corresponding fixed point $\hat{p}(t)$. This completes the proof of Theorem 1.

9. Trace monodromy and an isotopy principle

In this section we show the isotopy class assumption in Theorem 2 is important and necessary. To do this we first give a discussion of a topological obstruction to the extension of a holomorphic motion of a subset in the Riemann sphere parametrized by a non-simply connected Riemann surface. This discussion is first appeared in the paper [3]. Using this discussion we give a counter-example of a holomorphic motion of a finite subset in the Riemann sphere parametrized by any non-simply connected bounded domain in the complex plane that cannot be extended to a continuous (thus, a holomorphic) motion of the whole Riemann sphere parametrized by the same domain.

The topological obstruction is called *trace monodromy* in [3]. Suppose X is any hyperbolic Riemann surface, E a subset of $\overline{\mathbb{C}}$ containing $0, 1$ and ∞ and $card(E) \geq 4$. If h is a continuous motion of $X \times E \to \overline{\mathbb{C}}$, then for any $z \neq 0, 1, \infty \in E$, $h(t, z) = h^z(t) : X \to \mathbb{C}_{0,1} = \overline{\mathbb{C}} \setminus \{0, 1, \infty\}$ is continuous and induces a homomorphism

$$h^z_* : \pi_1(X) \to \pi_1(\mathbb{C}_{0,1}).$$

This is called the trace monodromy induced by h. By definition the *trace monodromy* is *trivial* if it maps every element of $\pi_1(X)$ to the identity of $\pi_1(\mathbb{C}_{0,1})$. We have the following result in [3], where a more general result is proved.

Theorem 7. *Suppose X is a Riemann surface with a basepoint t_0. Let $h : X \times E \to \overline{\mathbb{C}}$ be a normalized holomorphic motion of a finite set E with $card(E) \geq 4$ with parameter space X. If h can be extended to a quasiconformal motion of $\overline{\mathbb{C}}$ with the same parameter space by X, then the trace monodromy $h^z_* : \pi_1(X, t_0) \to \pi_1(\mathbb{C}_{0,1})$ for $z \neq 0, 1, \infty \in E$ is trivial.*

Proof. For the benefit of the reader, we outline the proof here. Suppose $H(t, z) : X \times \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ is a continuous motion which extends $h(t, z) : X \times E \to \overline{\mathbb{C}}$. Let $\gamma(s)$, $0 \leq s \leq 1$, be a simple closed curve in X with $\gamma(0) = \gamma(1) = t_0$. Let $\mu(s)$ be the Beltrami coefficient of $H(\gamma(s), \cdot)$ which is continuous on s. Then $H(\gamma(s), \cdot) = w^{\mu(s)}(\cdot)$ since both are quasiconformal maps fixing $0, 1, \infty$ with the same Beltrami coefficient. Let $K(s, u) = w^{u\mu(s)}(z) : [0, 1] \times [0, 1] \to$

$\mathbb{C}_{0,1}$ for any $z \neq 0, 1, \infty \in E$. Then it is a continuous map such that $K(s,1) = H(\gamma(s), z)$ and $K(s,0) = z$. Thus $H(\gamma(s), z)$ is a continuous curve in $\mathbb{C}_{0,1}$ homotopic to a point z in $\mathbb{C}_{0,1}$. This implies that the trace monodromy is trivial. □

Using this theorem, we construct a counter-example of a holomorphic motion of a finite subset of the Riemann sphere parametrized by any non-simply connected planar domain which does not satisfy our isotopy class assumption in Theorem 2.

Example 1. *Suppose X is a planar domain in the Riemann sphere $\overline{\mathbb{C}}$ such that $\overline{\mathbb{C}} - X$ has more than one connected components and at least one component contains more than one point. Let t_0 be the base point of X. Then for any finite subset E in $\overline{\mathbb{C}}$ with $\#(E) \geq 4$, there is a holomorphic motion $h(t,z) : X \times E \to \overline{\mathbb{C}}$ that cannot be extended to a quasiconformal motion of $\overline{\mathbb{C}}$ with parameter space X.*

Proof. Since $\overline{\mathbb{C}} - X$ has a component containing more than one point, we can use two points in this component and a square root map to map X into a half-plane. Then applying a Möbius transformation, we can assume that X is a bounded planar domain such that $\overline{\mathbb{C}} - X$ has a unbounded component and a bounded component.

Suppose $z_0 \neq 0, 1, \infty \in E$. Then we have a domain $\widetilde{X}$ and a conformal map $z = \phi(t) : X \to \widetilde{X}$ such that $\phi(t_0) = z_0$ and $\widetilde{X} \cap (E - \{z_0\}) = \emptyset$ and 0 is in a bounded component of $\overline{\mathbb{C}} - \widetilde{X}$ and $E - \{0, z_0\}$ are all in the unbounded component of $\overline{\mathbb{C}} - \widetilde{X}$. Define $h(t,z) = z$ for any $z \neq z_0$ and $t \in X$ and $\phi(t, z_0) = \phi(t)$. Then it is a holomorphic motion but the trace mondromy is not trivial. Thus from Theorem 7, it cannot be extended to a quasiconformal motion to the whole Riemann sphere parametrized by X. □

Remark 1. When $\#(E) = 4$ and $X = \mathbb{C} - \{0, 1\}$ is the trice-punctured sphere with a basepoint t_0, Douady constructed the following counter-example. Let $E = \{0, 1, \infty, t_0\}$, let $h(t,z) : X \times E \to \overline{\mathbb{C}}$ and define h by $h(t, 0) = 0$, $h(t, 1) = 1$, and $h(t, \infty) = \infty$, and $h(t, t_0) = t$. Then Douady showed that h is a maximal holomorphic motion and, therefore, cannot be extended further. Since an annulus A can be thought as a covering space of the thrice-punctured sphere, there is a covering map $\pi : A \to X$. Earle considered $\widetilde{h}(t,z) = (\pi^* h)(t,z) = h(\pi(t), z) : A \times E \to \overline{\mathbb{C}}$ and showed it is also a maximal holomorphic motion. Thus, it also cannot extended further. See [7] for these two counter-examples and the definition of a maximal holomorphic motion. The topological obstruction defined in [3] gives us more flexibility to construct more counter-examples. One can find other counter-examples when the parameter space is the punctured disk in [3] or an annulus. The counter-example given in Example 1 is for any finite subset with any non-simply connected planar domain.

Remark 2. In [3] it is shown that when $card(E) = 4$ a holomorphic motion $h : X \times E \to \overline{\mathbb{C}}$ can be extended to a holomorphic motion $\widetilde{h} : X \times \overline{\mathbb{C}} \to \overline{\mathbb{C}}$ if, and only if, its trace monodromy is trivial.

References

[1] L. V. Ahlfors. Lectures on Quasiconformal Mapping. volume 38 of University Lecture Series., Amer. Math. Soc., 2006.

[2] L. V. Ahlfors and G. Weill. A uniqueness theorem for Beltrami equations. Proc. Am. Math. Soc., Vol. 13:975-978, 1962.

[3] M. Beck, Y. Jiang, S. Mitra, and H. Shiga. Extending holomorphic motions and monodromy. Ann. Acad. Sci. Fenn. Math., Vol. 37:53-67, 2012.

[4] L. Bers. A non-standard integral equation with applications to quasiconformal mapping. Acta Math., 116:113-134, 1966.

[5] L. Bers and H. Royden. Holomorphic families of injections. Acta Math., 157:259-286, 1986.

[6] A. Douady and C. J. Earle. Conformally natural extension of homeomorhisms of the circle. Acta Math., 157:23-48, 1986.

[7] C. J. Earle. Some maximal holomorphic motions. In Lipa's Legacy: Proceedings of the Bers Colloquium. AMS Contemp. Math. Ser. **211**, 1997.

[8] C. J. Earle and C. T. McMullen. Quasiconformal isotopies. Holomorphic Functions and Moduli, I:143-154, 1988.

[9] F. P. Gardiner. The existence of Jenkins-Strebel differentials from Teichmüller theory. Amer. J. of Math., 99:1097-1104, 1975.

[10] F. P. Gardiner. Schiffer's interior variation and quasiconformal mappings. Duke Math. J., 42:371-380, 1975.

[11] F. P. Gardiner. Measured foliations and the minimal norm property for quadratic differentials. Acta Math., 152:57-76, 1984.

[12] F. P. Gardiner. Teichmüller Theory and Quadratic Differentials. John Wiley & Sons, New York, 1987.

[13] F. P. Gardiner and N. Lakic. Quasiconformal Teichmüller Theory. AMS, Providence, Rhode Island, 2000.

[14] F. P. Gardiner and H. Masur. Extremal length geometry of Teichmüller space. Complex Variables, 16:209-237, 1991.

[15] J. Hubbard and H. Masur. Quadratic differentials and foliatons. Acta Math., 142:221-274, 1979.

[16] J. A. Jenkins. On the existence of certain general extremal metrics. Ann. Math., 66:440-453, 1957.

[17] Y. Jiang and S. Mitra. Douady-earle section, holomorphic motions, and some applications. Contemporary Mathematics, AMS, 575:219-252, 2012.

[18] A. Leutbecher. Über Spitzen diskontinuierlichen Gruppen von lineargebrochenen transformationen. Math. Z., 100:183-200, 1967.

[19] R. Màñé, P. Sad, and D. Sullivan. On the dynamics of rational maps. Ann. Èc. Norm. Sup., 96:193-217, 1983.

[20] S. Nag. The Torelli spaces of punctured tori and spheres. Duke Math. J., 48:359-388, 1981.

[21] H. Renelt. Konstruktion gewisser quadratischer Differential mit von Dirichletintergralen. Math. Nachr., 73:125-142, 1976.

[22] H. Shimizu. On discontinuous groups operating on the product of upper half-planes. Ann. Math., 77:33-77, 1959.

[23] Z. Slodkowski. Holomorphic motions and polynomial hulls. Proc. Amer. Math. Soc., 111:347-355, 1991.

[24] K. Strebel. Quadratic Differentials. Springer-Verlag, Berlin & New York, 1984.

[25] D. P. Sullivan and W. P. Thurston. Extending holomorphic motions. Acta Math., 157:243-257, 1986.

Frederick P. Gardiner, Department of Mathematics, Brooklyn College, Brooklyn, NY 11210 and Department of Mathematics, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016. Email:frederick.gardiner@gmail.com

Yunping Jiang, Department of Mathematices, Queens College CUNY, Flushing, NY 11367 and Department of Mathematics, CUNY Graduate Center, 365 Fifth Avenue, New York, NY 10016. Email: yunping.jiang@qc.cuny.edu